



Emily Riehl

Johns Hopkins University

Do we need a new foundation for higher structures?

joint with Nikolai Kudasov and Jonathan Weinberger*



*also: Abdelrahman Aly Abouneqm, Fredrik Bakke,
César Bardomiano Martínez, Jonathan Campbell,
Matthias Hutzler, Kenji Maillard,
David Martínez Carpena, Nima Rasekh,
Florrie Verity, Tashi Walde

Plan



1. Computer formalization of mathematics
2. A computer proof assistant for higher category theory?
3. The **Rzk** proof assistant for simplicial homotopy type theory
4. Synthetic ∞ -category theory
5. A formalized proof of the ∞ -categorical Yoneda lemma



1

Computer formalization of mathematics



CAHIERS DE TOPOLOGIE
ET GÉOMÉTRIE DIFFÉRENTIELLE
CATÉGORIQUES

VOL. XXXII-1 (1991)

∞ -GROUPOIDS AND HOMOTOPY TYPES

by M.M. KAPRANOV and V.A. VOEVODSKY

RÉSUMÉ. Nous présentons une description de la catégorie homotopique des CW-complexes en termes des ∞ -groupoïdes. La possibilité d'une telle description a été suggérée par A. Grothendieck dans son mémoire "A la poursuite des champs".

It is well-known [GZ] that CW-complexes X such that $\pi_i(X, x) = 0$ for all $i \geq 2$, $x \in X$, are described, at the homotopy level, by groupoids. A. Grothendieck suggested, in his unpublished memoir [Gr], that this connection should have a higher-dimensional generalisation involving polycategories, viz. polycategorical analogues of groupoids. It is the purpose of this paper to establish such a generalisation.

- 15 statements =
4 theorems
+ 9 propositions
+ 1 lemma
+ 1 corollary
- 5 short “obvious” proofs + 3 proofs



CAHIERS DE TOPOLOGIE
ET GÉOMÉTRIE DIFFÉRENTIELLE
CATÉGORIQUES

VOL. XXXII-1 (1991)

∞ -GROUPOIDS AND HOMOTOPY TYPES

by M.M. KAPRANOV and V.A. VOEVODSKY

RÉSUMÉ. Nous présentons une description de la catégorie homotopique des CW-complexes en termes des ∞ -groupoïdes. La possibilité d'une telle description a été suggérée par A. Grothendieck dans son mémoire "A la poursuite des champs".

It is well-known [GZ] that CW-complexes X such that $\pi_i(X, x) = 0$ for all $i \geq 2$, $x \in X$, are described, at the homotopy level, by groupoids. A. Grothendieck suggested, in his unpublished memoir [Gr], that this connection should have a higher-dimensional generalisation involving polycategories, viz. polycategorical analogues of groupoids. It is the purpose of this paper to establish such a generalisation.

- Carlos Simpson's "Homotopy types of strict 3-groupoids" (1998) shows that the 3-type of S^2 can't be realized by a strict 3-groupoid — contradicting the last corollary.

- 15 statements =
4 theorems
+ 9 propositions
+ 1 lemma
+ 1 corollary
- 5 short "obvious" proofs + 3 proofs



CAHIERS DE TOPOLOGIE
ET GÉOMÉTRIE DIFFÉRENTIELLE
CATÉGORIQUES

VOL. XXXII-1 (1991)

∞ -GROUPOIDS AND HOMOTOPY TYPES

by M.M. KAPRANOV and V.A. VOEVODSKY

RÉSUMÉ. Nous présentons une description de la catégorie homotopique des CW-complexes en termes des ∞ -groupoïdes. La possibilité d'une telle description a été suggérée par A. Grothendieck dans son mémoire "A la poursuite des champs".

It is well-known [GZ] that CW-complexes X such that $\pi_i(X, x) = 0$ for all $i \geq 2$, $x \in X$, are described, at the homotopy level, by groupoids. A. Grothendieck suggested, in his unpublished memoir [Gr], that this connection should have a higher-dimensional generalisation involving polycategories, viz. polycategorical analogues of groupoids. It is the purpose of this paper to establish such a generalisation.

- Carlos Simpson's "Homotopy types of strict 3-groupoids" (1998) shows that the 3-type of S^2 can't be realized by a strict 3-groupoid — contradicting the last corollary.
- But no explicit mistake was found. Voevodsky: "I was sure that we were right until the fall of 2013 (!!)"

- 15 statements =
4 theorems
+ 9 propositions
+ 1 lemma
+ 1 corollary
- 5 short "obvious" proofs + 3 proofs



MATHEMATICS

The Origins and Motivations of Univalent Foundations

*A Personal Mission to Develop Computer Proof
Verification to Avoid Mathematical Mistakes*

By Vladimir Voevodsky • Published 2014

“A technical argument by a trusted author, which is hard to check and looks similar to arguments known to be correct, is hardly ever checked in detail.”

Computer formalized mathematics



Formalized mathematics, in tandem with other forms of computerized mathematics¹, provides better management of mathematical knowledge, an opportunity to carry out ever more complex and larger projects, and hitherto unseen levels of precision.

*— Andrej Bauer, “The dawn of formalized mathematics,”
delivered at the 8th European Congress of Mathematics*

¹Jacques Carette, William M. Farmer, Michael Kohlhase, and Florian Rabe. Big math and the one-brain barrier — the tetrapod model of mathematical knowledge. *Mathematical Intelligencer*, 43(1):78–87, 2021.

Computer formalized mathematics



Formalized mathematics, in tandem with other forms of computerized mathematics¹, provides better management of mathematical knowledge, an opportunity to carry out ever more complex and larger projects, and hitherto unseen levels of precision.

*— Andrej Bauer, “The dawn of formalized mathematics,”
delivered at the 8th European Congress of Mathematics*

Recent successes include:

- the **Kepler conjecture**, resolving a 1611 conjecture, 2003–2014, **HOL LIGHT**
- the **Feit-Thompson Odd Order Theorem**, a foundational result in the classification of finite simple groups, 2006–2012, **Coq**
- the **liquid tensor experiment**, formalizing condensed mathematics, 2020–2022, **LEAN**
- the **Brunerie number**, computing $\pi_4 S^3 \cong \mathbb{Z}/2\mathbb{Z}$, 2015–2022, **CUBICAL AGDA**

¹Jacques Carette, William M. Farmer, Michael Kohlhase, and Florian Rabe. Big math and the one-brain barrier — the tetrapod model of mathematical knowledge. *Mathematical Intelligencer*, 43(1):78–87, 2021.



2

A computer proof assistant for higher category
theory?

Rebuilding the pragmatic foundations for higher structures



I am pretty strongly convinced that there is an ongoing reversal in the collective consciousness of mathematicians: the homotopical picture of the world becomes the basic intuition, and if you want to get a discrete set, then you pass to the set of connected components of a space defined only up to homotopy ... Cantor's problems of the infinite recede to the background: from the very start, our images are so infinite that if you want to make something finite out of them, you must divide them by another infinity.

— Yuri Manin “We do not choose mathematics as our profession, it chooses us: Interview with Yuri Manin” by Mikhail Gelfand

∞ -categories in set theory



Essentially, ∞ -categories are 1-categories in which all the **sets** have been replaced by **∞ -groupoids** aka **homotopy types**:

sets :: ∞ -groupoids
categories :: ∞ -categories

∞ -categories in set theory



Essentially, ∞ -categories are 1-categories in which all the **sets** have been replaced by **∞ -groupoids** aka **homotopy types**:

sets :: ∞ -groupoids
categories :: ∞ -categories

Where

- categories have sets of objects, ∞ -categories have ∞ -groupoids of objects, and
- categories have hom-sets, ∞ -categories have ∞ -groupoidal mapping spaces.

∞ -categories in set theory



Essentially, ∞ -categories are 1-categories in which all the **sets** have been replaced by **∞ -groupoids** aka **homotopy types**:

sets :: ∞ -groupoids
categories :: ∞ -categories

Where

- categories have sets of objects, ∞ -categories have ∞ -groupoids of objects, and
- categories have hom-sets, ∞ -categories have ∞ -groupoidal mapping spaces.

While the axioms that turn a directed graph into a category are expressed in the language of set theory — a category has a composition function satisfying axioms expressed in first-order logic with equality

∞ -categories in set theory



Essentially, ∞ -categories are 1-categories in which all the **sets** have been replaced by **∞ -groupoids** aka **homotopy types**:

sets :: ∞ -groupoids
categories :: ∞ -categories

Where

- categories have sets of objects, ∞ -categories have ∞ -groupoids of objects, and
- categories have hom-sets, ∞ -categories have ∞ -groupoidal mapping spaces.

While the axioms that turn a directed graph into a category are expressed in the language of set theory — a category has a composition function satisfying axioms expressed in first-order logic with equality — composition in an ∞ -category, as a morphism between ∞ -groupoids, isn't a “function” in the traditional sense (since homotopy types do not have underlying sets of points).

∞ -categories in set theory



Essentially, ∞ -categories are 1-categories in which all the **sets** have been replaced by **∞ -groupoids** aka **homotopy types**:

sets :: ∞ -groupoids
categories :: ∞ -categories

Where

- categories have sets of objects, ∞ -categories have ∞ -groupoids of objects, and
- categories have hom-sets, ∞ -categories have ∞ -groupoidal mapping spaces.

While the axioms that turn a directed graph into a category are expressed in the language of set theory — a category has a composition function satisfying axioms expressed in first-order logic with equality — composition in an ∞ -category, as a morphism between ∞ -groupoids, isn't a “function” in the traditional sense (since homotopy types do not have underlying sets of points).


This is why ∞ -categories are so difficult to model within set theory.

Composing paths



In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”


of a space X , composites of paths are witnessed by higher paths:



Composing paths

In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”


of a space X , composites of paths are witnessed by higher paths:



Composing paths

In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”

of a space X , composites of paths are witnessed by higher paths:




Theorem. The space of composites of two paths f and g in X is contractible.

Composing paths

In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”

of a space X , composites of paths are witnessed by higher paths:



Theorem. The space of composites of two paths f and g in X is contractible.


Proof: The **space of composites** of paths f and g in X is defined by the pullback:

$$\begin{array}{ccc} \mathrm{Comp}(f, g) & \hookrightarrow & \mathrm{Map}(\Delta, X) \\ \downarrow & \lrcorner & \downarrow \text{restrict} \\ * & \xrightarrow{f \wedge g} & \mathrm{Map}(\Lambda, X) \end{array}$$

Composing paths

In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”

of a space X , composites of paths are witnessed by higher paths:



Theorem. The space of composites of two paths f and g in X is contractible.

Proof: The **space of composites** of paths f and g in X is defined by the pullback:


$$\begin{array}{ccccc}
 S^{n-1} & \longrightarrow & \text{Comp}(f, g) & \hookrightarrow & \text{Map}(\Delta, X) \\
 \downarrow & \nearrow \text{dashed} & \downarrow & \lrcorner & \downarrow \text{restrict} \\
 D^n & \longrightarrow & * & \xrightarrow{f \wedge g} & \text{Map}(\Lambda, X)
 \end{array}$$

A space is **contractible** just when any sphere S^{n-1} can be filled to a disk D^n for $n \geq 0$.

Composing paths

In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”

of a space X , composites of paths are witnessed by higher paths:



Theorem. The space of composites of two paths f and g in X is contractible.

Proof: The **space of composites** of paths f and g in X is defined by the pullback:


$$\begin{array}{ccccc}
 S^{n-1} & \longrightarrow & \text{Comp}(f, g) & \hookrightarrow & \text{Map}(\Delta, X) \\
 \downarrow & \nearrow & \downarrow & \nearrow & \downarrow \text{restrict} \\
 D^n & \longrightarrow & * & \xrightarrow{f \wedge g} & \text{Map}(\Lambda, X)
 \end{array}$$

A space is **contractible** just when any sphere S^{n-1} can be filled to a disk D^n for $n \geq 0$.

Composing paths

In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”

of a space X , composites of paths are witnessed by higher paths:



Theorem. The space of composites of two paths f and g in X is contractible.

Proof: The **space of composites** of paths f and g in X is defined by the pullback:


$$\begin{array}{ccc}
 S^{n-1} & \longrightarrow & \text{Comp}(f, g) \hookrightarrow \text{Map}(\Delta, X) \\
 \downarrow & \nearrow & \downarrow \text{restrict} \\
 D^n & \xrightarrow{\quad} & * \xrightarrow{f \wedge g} \text{Map}(\Lambda, X)
 \end{array}
 \quad \Leftrightarrow \quad
 \begin{array}{ccc}
 S^{n-1} \times \Delta \cup_{S^{n-1} \times \Lambda} D^n \times \Lambda & \longrightarrow & X \\
 \downarrow & \nearrow & \\
 D^n \times \Delta & &
 \end{array}$$

A space is **contractible** just when any sphere S^{n-1} can be filled to a disk D^n for $n \geq 0$.

Composing paths

In the **total singular complex** aka the **fundamental ∞ -groupoid** aka the **anima** or “soul”

of a space X , composites of paths are witnessed by higher paths:



Theorem. The space of composites of two paths f and g in X is contractible.

Proof: The **space of composites** of paths f and g in X is defined by the pullback:

$$\begin{array}{ccc}
 S^{n-1} & \longrightarrow & \text{Comp}(f, g) \hookrightarrow \text{Map}(\Delta, X) \\
 \downarrow & \nearrow & \downarrow \text{restrict} \\
 D^n & \xrightarrow{\quad} & * \xrightarrow{f \wedge g} \text{Map}(\Lambda, X)
 \end{array}
 \quad \Leftrightarrow \quad
 \begin{array}{ccc}
 S^{n-1} \times \Delta \cup_{S^{n-1} \times \Lambda} D^n \times \Lambda & \longrightarrow & X \\
 \downarrow & \nearrow & \\
 D^n \times \Delta & &
 \end{array}$$

A space is **contractible** just when any sphere S^{n-1} can be filled to a disk D^n for $n \geq 0$. The extension exists since the inclusion admits a continuous deformation retract. \square

Could ∞ -category theory be taught to undergraduates?

As far as we know, there are **no existing formalizations of ∞ -category theory** in any proof assistant library such as **LEAN-MATHLIB**, **AGDA-UNIMATH**, **COQ-HOTT**,...



Could ∞ -category theory be taught to undergraduates?



As far as we know, there are **no existing formalizations of ∞ -category theory** in any proof assistant library such as **LEAN-MATHLIB**, **AGDA-UNIMATH**, **COQ-HOTT**,...

Why not?

Could ∞ -category theory be taught to undergraduates?



As far as we know, there are **no existing formalizations of ∞ -category theory** in any proof assistant library such as **LEAN-MATHLIB**, **AGDA-UNIMATH**, **COQ-HoTT**,...

Why not?

Could ∞ -Category Theory Be Taught to Undergraduates?



Emily Riehl

1. The Algebra of Paths

It is natural to probe a suitably nice topological space X by means of its paths, the continuous functions from the standard unit interval $I = [0, 1] \subset \mathbb{R}$ to X . But what structure do the paths in X form?

To start, the paths form the edges of a directed graph whose vertices are the points of X : a path $p: I \rightarrow X$ defines an arrow from the point $p(0)$ to the point $p(1)$. Moreover,

this graph is reflexive, with the constant path rel_x at each point $x \in X$ defining a distinguished endomorphism.

Can this reflexive directed graph be given the structure of a category? To do so, it is natural to define the composite of a path p from x to y and a path q from y to z by gluing together these continuous maps—i.e., by concatenating the paths—and then by reparametrizing via the homeomorphism $I \cong I \cup_{[1,0]} I$ that traverses each path at double speed:

$$\begin{array}{c} I \xrightarrow{p} I \cup_{[1,0]} I \xrightarrow{q} X \\ \quad \quad \quad \text{cong} \end{array} \quad (1.1)$$

But the composition operation \circ fails to be associative or unital. In general, given a path r from z to u , the

The traditional foundations of mathematics are not really suitable for “higher mathematics” such as ∞ -category theory, where the basic objects are built out of higher-dimensional types instead of mere sets. However, there are proposals for new foundations for mathematics that are closer to mathematician’s core intuitions, based on Martin-Löf’s dependent type theory such as

- homotopy type theory,
- higher observational type theory, and the
- **simplicial type theory**, that we use here.

Emily Riehl is a professor of mathematics at Johns Hopkins University. Her email address is eriehl@jhu.edu.
Communicated by Notion Associate Editor Simon S. S. For permission to reprint this article, please contact: reprint-permission@ams.org.
DOI: <https://doi.org/10.1090/noti2022>

∞ -categories in homotopy type theory

The identity type family gives each type the structure of an ∞ -groupoid: each type A has a family of identity types over $x, y : A$ whose terms $p : x =_A y$ are called **paths**.



∞ -categories in homotopy type theory



The identity type family gives each type the structure of an ∞ -groupoid: each type A has a family of identity types over $x, y : A$ whose terms $p : x =_A y$ are called **paths**. In a “directed” extension of homotopy type theory introduced in

Emily Riehl and Michael Shulman, *A type theory for synthetic ∞ -categories*,
Higher Structures 1(1):116–193, 2017

each type A also has a family of hom types $\text{Hom}_A(x, y)$ over $x, y : A$ whose terms $f : \text{Hom}_A(x, y)$ are called **arrows**.

∞ -categories in homotopy type theory



The identity type family gives each type the structure of an ∞ -groupoid: each type A has a family of identity types over $x, y : A$ whose terms $p : x =_A y$ are called **paths**. In a “directed” extension of homotopy type theory introduced in

Emily Riehl and Michael Shulman, *A type theory for synthetic ∞ -categories*,
Higher Structures 1(1):116–193, 2017

each type A also has a family of hom types $\text{Hom}_A(x, y)$ over $x, y : A$ whose terms $f : \text{Hom}_A(x, y)$ are called **arrows**.

defn (Riehl–Shulman after Joyal and Rezk). A type A is an ∞ -category if:

∞ -categories in homotopy type theory



The identity type family gives each type the structure of an ∞ -groupoid: each type A has a family of identity types over $x, y : A$ whose terms $p : x =_A y$ are called **paths**. In a “directed” extension of homotopy type theory introduced in

Emily Riehl and Michael Shulman, *A type theory for synthetic ∞ -categories*,
Higher Structures 1(1):116–193, 2017

each type A also has a family of hom types $\text{Hom}_A(x, y)$ over $x, y : A$ whose terms $f : \text{Hom}_A(x, y)$ are called **arrows**.

defn (Riehl–Shulman after Joyal and Rezk). A type A is an ∞ -category if:

- Every pair of arrows $f : \text{Hom}_A(x, y)$ and $g : \text{Hom}_A(y, z)$ has a **unique composite**, defining a term $g \circ f : \text{Hom}_A(x, z)$.

∞ -categories in homotopy type theory



The identity type family gives each type the structure of an ∞ -groupoid: each type A has a family of identity types over $x, y : A$ whose terms $p : x =_A y$ are called **paths**. In a “directed” extension of homotopy type theory introduced in

Emily Riehl and Michael Shulman, *A type theory for synthetic ∞ -categories*,
Higher Structures 1(1):116–193, 2017

each type A also has a family of hom types $\text{Hom}_A(x, y)$ over $x, y : A$ whose terms $f : \text{Hom}_A(x, y)$ are called **arrows**.

defn (Riehl–Shulman after Joyal and Rezk). A type A is an ∞ -category if:

- Every pair of arrows $f : \text{Hom}_A(x, y)$ and $g : \text{Hom}_A(y, z)$ has a **unique composite**, defining a term $g \circ f : \text{Hom}_A(x, z)$.
- Paths in A are equivalent to **isomorphisms** in A .

∞ -categories in homotopy type theory



The identity type family gives each type the structure of an ∞ -groupoid: each type A has a family of identity types over $x, y : A$ whose terms $p : x =_A y$ are called **paths**. In a “directed” extension of homotopy type theory introduced in

Emily Riehl and Michael Shulman, *A type theory for synthetic ∞ -categories*,
Higher Structures 1(1):116–193, 2017

each type A also has a family of hom types $\text{Hom}_A(x, y)$ over $x, y : A$ whose terms $f : \text{Hom}_A(x, y)$ are called **arrows**.

defn (Riehl–Shulman after Joyal and Rezk). A type A is an ∞ -category if:

- Every pair of arrows $f : \text{Hom}_A(x, y)$ and $g : \text{Hom}_A(y, z)$ has a **unique composite**, defining a term $g \circ f : \text{Hom}_A(x, z)$.
- Paths in A are equivalent to **isomorphisms** in A .

With more of the work being done by the foundation system, perhaps someday ∞ -category theory will be easy enough to teach to undergraduates?



3

The **RZK** proof assistant for simplicial homotopy
type theory

Simplicial homotopy type theory



In **simplicial type theory**, types may depend on other types and also on **shapes**, which are polytopes $\Phi := \{\vec{t} : \mathcal{Z}^n \mid \phi(\vec{t})\}$ cut out of a directed cube by a formula $\phi(\vec{t})$ called a **tope**.

Simplicial homotopy type theory



In **simplicial type theory**, types may depend on other types and also on **shapes**, which are polytopes $\Phi := \{\vec{t} : \mathcal{Z}^n \mid \phi(\vec{t})\}$ cut out of a directed cube by a formula $\phi(\vec{t})$ called a **tope**.

- **Shapes** and their defining **topes** are described syntactically in a language using the symbols $\top, \perp, \wedge, \vee, \equiv$ and $0, 1, \leq$ satisfying **intuitionistic logic** and **strict interval axioms**:
e.g., $\Delta^n := \{(t_1, \dots, t_n) : \mathcal{Z}^n \mid t_n \leq \dots \leq t_1\}$.
- The shape defined by $\phi \vee \psi$ is the **strict pushout** of the shapes defined by ϕ and ψ over $\phi \wedge \psi$: e.g., $\partial\Delta^1 := \{t : \mathcal{Z} \mid (t \equiv 0) \vee (t \equiv 1)\}$ is the coproduct of two points.
- **Shape inclusions** $\Phi \subset \Psi$ arise from implications in intuitionistic logic: e.g., the topes

$$\Delta^2 := \{(t_1, t_2) : \mathcal{Z}^2 \mid t_2 \leq t_1\}$$

$$\partial\Delta^2 := \{(t_1, t_2) : \mathcal{Z}^2 \mid (t_2 \leq t_1) \wedge ((0 \equiv t_2) \vee (t_2 \equiv t_1) \vee (t_1 \equiv 1))\}$$

$$\Lambda_1^2 := \{(t_1, t_2) : \mathcal{Z}^2 \mid (t_2 \leq t_1) \wedge ((0 \equiv t_2) \vee (t_1 \equiv 1))\}$$

define shape inclusions $\Lambda_1^2 \subset \partial\Delta^2 \subset \Delta^2$.

Extension types



Formation rule for extension types

$$\frac{\Phi \subset \Psi \text{ shape} \quad A \text{ type} \quad a : \Phi \rightarrow A}{\left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \downarrow & & \uparrow \\ \Psi & \xrightarrow{\quad} & \end{array} \right\rangle \text{ type}}$$

Extension types



Formation rule for extension types

$$\frac{\Phi \subset \Psi \text{ shape} \quad A \text{ type} \quad a : \Phi \rightarrow A}{\left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \downarrow & \nearrow & \\ \Psi & & \end{array} \right\rangle \text{ type}}$$

A term $f : \left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \downarrow & \nearrow & \\ \Psi & & \end{array} \right\rangle$ defines

Extension types



Formation rule for extension types

$$\frac{\Phi \subset \Psi \text{ shape} \quad A \text{ type} \quad a : \Phi \rightarrow A}{\left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \Downarrow & \nearrow & \\ \Psi & & \end{array} \right\rangle \text{ type}}$$

A term $f : \left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \Downarrow & \nearrow & \\ \Psi & & \end{array} \right\rangle$ defines

$f : \Psi \rightarrow A$ so that $f(t) \equiv a(t)$ for $t : \Phi$.

Extension types



Formation rule for extension types

$$\frac{\Phi \subset \Psi \text{ shape} \quad A \text{ type} \quad a : \Phi \rightarrow A}{\left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \downarrow & \nearrow & \\ \Psi & & \end{array} \right\rangle \text{ type}}$$

A term $f : \left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \downarrow & \nearrow & \\ \Psi & & \end{array} \right\rangle$ defines

$f : \Psi \rightarrow A$ so that $f(t) \equiv a(t)$ for $t : \Phi$.

The simplicial type theory allows us to *prove* equivalences between extension types along composites or products of shape inclusions.

A formalized proof of the ∞ -categorical Yoneda lemma



Our initial aim was to write a formalized proof of the ∞ -categorical Yoneda lemma.

github.com/emilyriehl/yoneda or emilyriehl.github.io/yoneda/

- proof from Emily Riehl & Mike Shulman, *A type theory for synthetic ∞ -categories*, Higher Structures 2017.
- formalizations written by Nikolai Kudasov, Emily Riehl, Jonathan Weinberger.
- completed March 12 – April 17, 2023

A formalized proof of the ∞ -categorical Yoneda lemma



Our initial aim was to write a formalized proof of the ∞ -categorical Yoneda lemma.

github.com/emilyriehl/yoneda or emilyriehl.github.io/yoneda/

- proof from Emily Riehl & Mike Shulman, [A type theory for synthetic \$\infty\$ -categories](#), Higher Structures 2017.
- formalizations written by [Nikolai Kudasov](#), [Emily Riehl](#), [Jonathan Weinberger](#).
- completed March 12 – April 17, 2023

Another objective is to compare ∞ -category theory in simplicial type theory with ordinary category theory in traditional foundations. Thus,

- We've included a formalization of the 1-categorical Yoneda lemma in Lean by [Sina Hazratpour](#) as part of an Introduction to Proofs course at Johns Hopkins.
- We wrote a first version of [yoneda-lemma-precategories.lagda.md](#).

A formalized proof of the ∞ -categorical Yoneda lemma



Our initial aim was to write a formalized proof of the ∞ -categorical Yoneda lemma.

github.com/emilyriehl/yoneda or emilyriehl.github.io/yoneda/

- proof from Emily Riehl & Mike Shulman, [A type theory for synthetic \$\infty\$ -categories](#), Higher Structures 2017.
- formalizations written by [Nikolai Kudasov](#), [Emily Riehl](#), [Jonathan Weinberger](#).
- completed March 12 – April 17, 2023

Another objective is to compare ∞ -category theory in simplicial type theory with ordinary category theory in traditional foundations. Thus,

- We've included a formalization of the 1-categorical Yoneda lemma in Lean by [Sina Hazratpour](#) as part of an Introduction to Proofs course at Johns Hopkins.
- We wrote a first version of [yoneda-lemma-precategories.lagda.md](#).

More recently, we've professionalized our library, implementing a style guide suggested by [Fredrik Bakke](#), and invited new contributors to a broader project of formalizing synthetic ∞ -category theory:

github.com/rzk-lang/sHoTT or rzk-lang.github.io/sHoTT



4

Synthetic ∞ -category theory

Hom types



In the simplicial type theory, any type A has a family of **hom types** depending on two terms in $x, y : A$:

$$\mathbf{Hom}_A(x, y) := \left\langle \begin{array}{ccc} \partial\Delta^1 & \xrightarrow{[x,y]} & A \\ \Downarrow & \nearrow & \\ \Delta^1 & & \end{array} \right\rangle \text{ type}$$

A term $f : \mathbf{Hom}_A(x, y)$ defines an **arrow** in A from x to y .

We think of the type $\mathbf{Hom}_A(x, y)$ as the **mapping space** in A from x to y .

Hom types



In the simplicial type theory, any type A has a family of **hom types** depending on two terms in $x, y : A$:

$$\mathbf{Hom}_A(x, y) := \left\langle \begin{array}{ccc} \partial\Delta^1 & \xrightarrow{[x,y]} & A \\ \Downarrow & \nearrow & \\ \Delta^1 & & \end{array} \right\rangle \text{ type}$$

A term $f : \mathbf{Hom}_A(x, y)$ defines an **arrow** in A from x to y .

We think of the type $\mathbf{Hom}_A(x, y)$ as the **mapping space** in A from x to y .

A type A also has a family of **identity types** or **path spaces** $x = y$ depending on two terms in $x, y : A$, which we will connect to the hom-types momentarily.



defn (Riehl–Shulman after Joyal). A type A is a **pre- ∞ -category** if every pair of arrows $f : \mathbf{Hom}_A(x, y)$ and $g : \mathbf{Hom}_A(y, z)$ has a **unique composite**, i.e.,

$$\left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[f,g]} & A \\ \Downarrow & \nearrow \text{dashed} & \\ \Delta^2 & & \end{array} \right\rangle \quad \text{is contractible.}^a$$

^aA type C is contractible just when $\sum_{c:C} \prod_{x:C} c = x$.



defn (Riehl–Shulman after Joyal). A type A is a **pre- ∞ -category** if every pair of arrows $f : \mathbf{Hom}_A(x, y)$ and $g : \mathbf{Hom}_A(y, z)$ has a **unique composite**, i.e.,

$$\left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[f,g]} & A \\ \Downarrow & \nearrow \text{dashed} & \\ \Delta^2 & & \end{array} \right\rangle \quad \text{is contractible.}^a$$

^aA type C is contractible just when $\sum_{c:C} \prod_{x:C} c = x$.

By contractibility, $\left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[f,g]} & A \\ \Downarrow & \nearrow \text{dashed} & \\ \Delta^2 & & \end{array} \right\rangle$ has a unique inhabitant $\mathbf{comp}_{f,g} : \Delta^2 \rightarrow A$.

Write $g \circ f : \mathbf{Hom}_A(x, z)$ for its inner face, *the* composite of f and g .

Identity arrows



For any $x : A$, the constant function defines a term

$$\text{id}_x := \lambda t.x : \text{Hom}_A(x, x) := \left\langle \begin{array}{ccc} \partial\Delta^1 & \xrightarrow{[x,x]} & A \\ \Downarrow & \nearrow & \\ \Delta^1 & & \end{array} \right\rangle,$$

which we denote by id_x and call the **identity arrow**.

Identity arrows



For any $x : A$, the constant function defines a term

$$\text{id}_x := \lambda t. x : \text{Hom}_A(x, x) := \left\langle \begin{array}{ccc} \partial \Delta^1 & \xrightarrow{[x, x]} & A \\ \Downarrow & \nearrow & \\ \Delta^1 & & \end{array} \right\rangle,$$

which we denote by id_x and call the **identity arrow**.

For any $f : \text{Hom}_A(x, y)$ in a pre- ∞ -category A , the term in the contractible type

$$\lambda(s, t). f(t) : \left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[\text{id}_x, f]} & A \\ \Downarrow & \nearrow & \\ \Delta^2 & & \end{array} \right\rangle$$

witnesses the unit axiom $f = f \circ \text{id}_x$.

Associativity of composition




Prop. In a pre- ∞ -category A , composition is associative: for any arrows $f : \mathbf{Hom}_A(x, y)$, $g : \mathbf{Hom}_A(y, z)$, and $h : \mathbf{Hom}_A(z, w)$, we have $h \circ (g \circ f) = (h \circ g) \circ f$.

Associativity of composition



Prop. In a pre- ∞ -category A , composition is associative: for any arrows $f : \text{Hom}_A(x, y)$, $g : \text{Hom}_A(y, z)$, and $h : \text{Hom}_A(z, w)$, we have $h \circ (g \circ f) = (h \circ g) \circ f$.

Proof: Consider the composable arrows in the pre- ∞ -category $\Delta^1 \rightarrow A$:




Associativity of composition



Prop. In a pre- ∞ -category A , composition is associative: for any arrows $f : \text{Hom}_A(x, y)$, $g : \text{Hom}_A(y, z)$, and $h : \text{Hom}_A(z, w)$, we have $h \circ (g \circ f) = (h \circ g) \circ f$.

Proof: Consider the composable arrows in the pre- ∞ -category $\Delta^1 \rightarrow A$:




Composing defines a term in the type $\Delta^2 \rightarrow (\Delta^1 \rightarrow A)$

Associativity of composition



Prop. In a pre- ∞ -category A , composition is associative: for any arrows $f : \mathbf{Hom}_A(x, y)$, $g : \mathbf{Hom}_A(y, z)$, and $h : \mathbf{Hom}_A(z, w)$, we have $h \circ (g \circ f) = (h \circ g) \circ f$.

Proof: Consider the composable arrows in the pre- ∞ -category $\Delta^1 \rightarrow A$:



Composing defines a term in the type $\Delta^2 \rightarrow (\Delta^1 \rightarrow A)$ which defines an arrow $\ell : \mathbf{Hom}_A(x, w)$ so that $\ell = h \circ (g \circ f)$ and $\ell = (h \circ g) \circ f$.

Isomorphisms



An arrow $f: \text{Hom}_A(x, y)$ in a pre- ∞ -category is an **isomorphism** if it has a two-sided inverse $g: \text{Hom}_A(y, x)$. However, the type

$$\sum_{g: \text{Hom}_A(y, x)} (g \circ f = \text{id}_x) \times (f \circ g = \text{id}_y)$$

has higher-dimensional structure and is *not* a **proposition**.

Isomorphisms



An arrow $f: \text{Hom}_A(x, y)$ in a pre- ∞ -category is an **isomorphism** if it has a two-sided inverse $g: \text{Hom}_A(y, x)$. However, the type

$$\sum_{g: \text{Hom}_A(y, x)} (g \circ f = \text{id}_x) \times (f \circ g = \text{id}_y)$$

has higher-dimensional structure and is *not* a **proposition**. Instead define

$$\text{is-iso}(f) := \left(\sum_{g: \text{Hom}_A(y, x)} g \circ f = \text{id}_x \right) \times \left(\sum_{h: \text{Hom}_A(y, x)} f \circ h = \text{id}_y \right).$$

Isomorphisms



An arrow $f: \text{Hom}_A(x, y)$ in a pre- ∞ -category is an **isomorphism** if it has a two-sided inverse $g: \text{Hom}_A(y, x)$. However, the type

$$\sum_{g: \text{Hom}_A(y, x)} (g \circ f = \text{id}_x) \times (f \circ g = \text{id}_y)$$

has higher-dimensional structure and is *not* a **proposition**. Instead define

$$\text{is-iso}(f) := \left(\sum_{g: \text{Hom}_A(y, x)} g \circ f = \text{id}_x \right) \times \left(\sum_{h: \text{Hom}_A(y, x)} f \circ h = \text{id}_y \right).$$

For $x, y: A$, the **type of isomorphisms** from x to y is:

$$x \cong_A y := \sum_{f: \text{Hom}_A(x, y)} \text{is-iso}(f).$$

∞ -categories



By path induction, to define a map

$$\text{iso-eq} : (x =_A y) \rightarrow (x \cong_A y)$$

for all $x, y : A$ it suffices to define

$$\text{iso-eq}(\text{refl}_x) := \text{id}_x.$$

∞ -categories



By path induction, to define a map

$$\text{iso-eq} : (x =_A y) \rightarrow (x \cong_A y)$$

for all $x, y : A$ it suffices to define

$$\text{iso-eq}(\text{refl}_x) := \text{id}_x.$$

defn (Riehl–Shulman after Rezk). A pre- ∞ -category A is ∞ -category iff every isomorphism is an identity



By path induction, to define a map

$$\text{iso-eq} : (x =_A y) \rightarrow (x \cong_A y)$$

for all $x, y : A$ it suffices to define

$$\text{iso-eq}(\text{refl}_x) := \text{id}_x.$$

defn (Riehl–Shulman after Rezk). A pre- ∞ -category A is ∞ -category iff every isomorphism is an identity, i.e., iff the map

$$\text{iso-eq} : \prod_{x, y : A} (x =_A y) \rightarrow (x \cong_A y)$$

is an equivalence.

∞ -groupoids



Similarly by path induction define

$$\text{arr-eq} : (x =_A y) \rightarrow \text{Hom}_A(x, y)$$

for all $x, y : A$ by $\text{arr-eq}(\text{refl}_x) := \text{id}_x$.

∞ -groupoids



Similarly by path induction define

$$\text{arr-eq} : (x =_A y) \rightarrow \text{Hom}_A(x, y)$$

for all $x, y : A$ by $\text{arr-eq}(\text{refl}_x) := \text{id}_x$.

A type A is an ∞ -groupoid iff every arrow is an identity, i.e., iff arr-eq is an equivalence.

∞ -groupoids



Similarly by path induction define

$$\text{arr-eq} : (x =_A y) \rightarrow \text{Hom}_A(x, y)$$

for all $x, y : A$ by $\text{arr-eq}(\text{refl}_x) := \text{id}_x$.

A type A is an ∞ -groupoid iff every arrow is an identity, i.e., iff arr-eq is an equivalence.

Prop. A type is an ∞ -groupoid if and only if it is an ∞ -category and all of its arrows are isomorphisms.

Proof:

$$\begin{array}{ccc} x =_A y & \xrightarrow{\text{arr-eq}} & \text{Hom}_A(x, y) \\ & \searrow \text{iso-eq} & \nearrow \\ & x \cong_A y & \end{array}$$

∞ -categories for undergraduates



defn. An ∞ -groupoid is a type in which arrows are equivalent to identities:

$\text{arr-eq} : (x =_A y) \rightarrow \text{Hom}_A(x, y)$ is an equivalence.

∞ -categories for undergraduates



defn. An ∞ -groupoid is a type in which arrows are equivalent to identities:

$\text{arr-eq}: (x =_A y) \rightarrow \text{Hom}_A(x, y)$ is an equivalence.

defn. An ∞ -category is a type

- which has unique binary composites of arrows:

$$\left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[f,g]} & A \\ \Downarrow & \searrow \text{dashed} & \\ \Delta^2 & & \end{array} \right\rangle \quad \text{is contractible}$$

∞ -categories for undergraduates



defn. An ∞ -groupoid is a type in which arrows are equivalent to identities:

$\text{arr-eq}: (x =_A y) \rightarrow \text{Hom}_A(x, y)$ is an equivalence.

defn. An ∞ -category is a type

- which has unique binary composites of arrows:

$$\left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[f,g]} & A \\ \downarrow & \searrow & \\ \Delta^2 & & \end{array} \right\rangle \quad \text{is contractible}$$

- and in which isomorphisms are equivalent to identities:

$\text{iso-eq}: (x =_A y) \rightarrow (x \cong_A y)$ is an equivalence.



5

A formalized proof of the ∞ -categorical Yoneda lemma

Stating the Yoneda lemma



Let A be a pre- ∞ -category and fix $a, b : A$.

Yoneda lemma. Evaluation at the identity defines an equivalence

$$\text{evid} := \lambda \phi. \phi_a(\text{id}_a) : \left(\prod_{x:A} \text{Hom}_A(a, x) \rightarrow \text{Hom}_A(b, x) \right) \rightarrow \text{Hom}_A(b, a)$$

Stating the Yoneda lemma



Let A be a pre- ∞ -category and fix $a, b : A$.

Yoneda lemma. Evaluation at the identity defines an equivalence

$$\text{evid} := \lambda \phi. \phi_a(\text{id}_a) : \left(\prod_{x:A} \text{Hom}_A(a, x) \rightarrow \text{Hom}_A(b, x) \right) \rightarrow \text{Hom}_A(b, a)$$

While terms $\phi : \prod_{x:A} \text{Hom}_A(a, x) \rightarrow \text{Hom}_A(b, x)$ are just families of maps

$$\phi_x : \text{Hom}_A(a, x) \rightarrow \text{Hom}_A(b, x)$$

indexed by terms $x : A$ such families are automatically **natural**:

Prop. Any family of maps $\phi : \prod_{x:A} \text{hom}_A(a, x) \rightarrow \text{hom}_A(b, x)$ is **natural**:

for any $g : \text{hom}_A(a, y)$ and $h : \text{hom}_A(y, z)$

$$h \circ \phi_y(g) = \phi_z(h \circ g).$$

Proving the Yoneda lemma



Let A be a pre- ∞ -category and fix $a, b : A$.

Yoneda lemma. Evaluation at the identity defines an equivalence

$$\text{eval} := \lambda \phi. \phi_a(\text{id}_a) : \left(\prod_{x:A} \text{Hom}_A(a, x) \rightarrow \text{Hom}_A(b, x) \right) \rightarrow \text{Hom}_A(b, a)$$

The proof is (a simplification of) the standard argument for 1-categories!

Proving the Yoneda lemma



Let A be a pre- ∞ -category and fix $a, b : A$.

Yoneda lemma. Evaluation at the identity defines an equivalence

$$\text{evid} := \lambda \phi. \phi_a(\text{id}_a) : \left(\prod_{x:A} \text{Hom}_A(a, x) \rightarrow \text{Hom}_A(b, x) \right) \rightarrow \text{Hom}_A(b, a)$$

The proof is (a simplification of) the standard argument for 1-categories!

Proof: Define an inverse map by

$$\text{yon} := \lambda f. \lambda x. \lambda g. g \circ f : \text{Hom}_A(b, a) \rightarrow \left(\prod_{x:A} \text{Hom}_A(a, x) \rightarrow \text{Hom}_A(b, x) \right).$$

Proving the Yoneda lemma



Let A be a pre- ∞ -category and fix $a, b : A$.

Yoneda lemma. Evaluation at the identity defines an equivalence

$$\mathbf{eval} := \lambda \phi. \phi_a(\mathbf{id}_a) : \left(\prod_{x:A} \mathrm{Hom}_A(a, x) \rightarrow \mathrm{Hom}_A(b, x) \right) \rightarrow \mathrm{Hom}_A(b, a)$$

The proof is (a simplification of) the standard argument for 1-categories!

Proof: Define an inverse map by

$$\mathbf{yon} := \lambda f. \lambda x. \lambda g. g \circ f : \mathrm{Hom}_A(b, a) \rightarrow \left(\prod_{x:A} \mathrm{Hom}_A(a, x) \rightarrow \mathrm{Hom}_A(b, x) \right).$$

By definition, $\mathbf{eval} \circ \mathbf{yon}(f) := \mathbf{id}_a \circ f$, and since $\mathbf{id}_a \circ f = f$, so $\mathbf{eval} \circ \mathbf{yon}(f) = f$.

Proving the Yoneda lemma

Let A be a pre- ∞ -category and fix $a, b : A$.

Yoneda lemma. Evaluation at the identity defines an equivalence

$$\mathbf{eval} := \lambda \phi. \phi_a(\mathbf{id}_a) : \left(\prod_{x:A} \mathrm{Hom}_A(a, x) \rightarrow \mathrm{Hom}_A(b, x) \right) \rightarrow \mathrm{Hom}_A(b, a)$$

The proof is (a simplification of) the standard argument for 1-categories!

Proof: Define an inverse map by

$$\mathbf{yon} := \lambda f. \lambda x. \lambda g. g \circ f : \mathrm{Hom}_A(b, a) \rightarrow \left(\prod_{x:A} \mathrm{Hom}_A(a, x) \rightarrow \mathrm{Hom}_A(b, x) \right).$$

By definition, $\mathbf{eval} \circ \mathbf{yon}(f) := \mathbf{id}_a \circ f$, and since $\mathbf{id}_a \circ f = f$, so $\mathbf{eval} \circ \mathbf{yon}(f) = f$.
Similarly, by definition, $\mathbf{yon} \circ \mathbf{eval}(\phi)_x(g) := g \circ \phi_a(\mathbf{id}_a)$. By naturality of ϕ and another identity law $g \circ \phi_a(\mathbf{id}_a) = \phi_x(g \circ \mathbf{id}_a) = \phi_x(g)$, so $\mathbf{yon} \circ \mathbf{eval}(\phi)_x(g) = \phi_x(g)$. \square

Conclusions and future work

Observations:

- ∞ -category theory is significantly easier to formalize in a foundation system based on homotopy type theory.
- By moving much of the complexity of “higher structures” into the background foundation system, the gap between ∞ -category theory and 1-category narrows substantially.
- A computer proof assistant is a fantastic tool for learning to write proofs in new foundations — indeed, through formalization in **RZK** we caught an error of circular reasoning in the **Riehl–Shulman** paper!

Future work:

- We would love help formalizing more results from ∞ -category theory in **RZK**.
- But the initial version of the simplicial type theory is not sufficiently powerful to prove all results about ∞ -categories, so further extensions of this synthetic framework are needed.

References

- Emily Riehl, [Could \$\infty\$ -category theory be taught to undergraduates?](#), Notices of the AMS 70(5):727–736, May 2023; [arXiv:2302.07855](#)
- Nikolai Kudasov, Emily Riehl, Jonathan Weinberger, [Formalizing the \$\infty\$ -categorical Yoneda lemma](#), CPP 2024: 274–290; [arXiv:2309.08340](#)
- Emily Riehl and Michael Shulman, [A type theory for synthetic \$\infty\$ -categories](#), Higher Structures 1(1):116–193, 2017; [arXiv:1705.07442](#)
- César Barmann, [Limits and colimits of synthetic \$\infty\$ -categories](#), [arXiv:2202.12386](#)
- Ulrik Buchholtz, Jonathan Weinberger, [Synthetic fibered \$\(\infty, 1\)\$ -category theory](#), Higher Structures 7(1): 74–165, 2023; [arXiv:2105.01724](#)

Danke!