



Emily Riehl

Johns Hopkins University

# A synthetic theory of $\infty$ -categories in homotopy type theory

joint with Michael Shulman

Octoberfest, Carnegie Mellon University

# Motivation



Why do I study *category theory*?

— I find category theoretic arguments to be aesthetically appealing.

What draws me to *homotopy type theory*?

— I find homotopy type theoretic arguments to be aesthetically appealing.



1. Homotopy type theory
2. A type theory for synthetic  $(\infty, 1)$ -categories
3. Segal types and Rezk types
4. The synthetic theory of  $(\infty, 1)$ -categories

Main takeaway: the [dependent Yoneda lemma](#) is a directed analogue of [path induction](#) in HoTT.



# Homotopy type theory

# Types, terms, and type constructors



Homotopy type theory has:

- types  $A, B, \dots$
- terms  $x : A, y : B$
- dependent types  $x : A \vdash B(x)$  type,  $x, y : A \vdash B(x, y)$  type

Type constructors build new types and terms from given ones:

- products  $A \times B$ , coproducts  $A + B$ , function types  $A \rightarrow B$ ,
- dependent sums  $\sum_{x:A} B(x)$ , dependent products  $\prod_{x:A} B(x)$ , and identity types  $x, y : A \vdash x =_A y$ .

Propositions as types:

$A \times B$		$A$ and $B$	$\sum_{x:A} B(x)$		$\exists x. B(x)$
$A + B$		$A$ or $B$	$\prod_{x:A} B(x)$		$\forall x. B(x)$
$A \rightarrow B$		$A$ implies $B$	$x =_A y$		$x$ equals $y$

# Identity types



Formation and introduction rules for identity types

$$\frac{x, y : A}{x =_A y \text{ type}}$$

$$\frac{x : A}{\text{refl}_x : x =_A x}$$

$$\text{Semantics} \left\{ \begin{array}{l} \sum_{x,y:A} x =_A y \\ \downarrow \\ A \xrightarrow{\Delta} A \times A \end{array} \right.$$

$\lambda x. \text{refl}_x$  (indicated by a dashed orange arrow pointing from  $A$  to  $\sum_{x,y:A} x =_A y$ )

Hence  $\sum_{x,y:A} x =_A y$  is interpreted as the **path space** of  $A$  and a term  $p : x =_A y$  may be thought of as a **path** from  $x$  to  $y$  in  $A$ .

# Path induction



The identity type family is freely generated by the terms  $\text{refl}_x : x =_A x$ .

**Path induction:** If  $B(x, y, p)$  is a type family dependent on  $x, y : A$  and  $p : x =_A y$ , then there is a function

$$\text{path-ind} : \left( \prod_{x:A} B(x, x, \text{refl}_x) \right) \rightarrow \left( \prod_{x,y:A} \prod_{p:x=_A y} B(x, y, p) \right).$$

Thus, to prove  $B(x, y, p)$  it suffices to assume  $y$  is  $x$  and  $p$  is  $\text{refl}_x$ .

The  $\infty$ -groupoid structure of  $A$  with

- terms  $x : A$  as objects
- paths  $p : x =_A y$  as 1-morphisms
- paths of paths  $h : p =_{x=_A y} q$  as 2-morphisms, ...

arises automatically from the path induction principle.




2

A type theory for synthetic  
 $(\infty, 1)$ -categories



# The intended model



**Theorem** ([Shulman](#)). Homotopy type theory is modeled by the category of [Reedy](#) fibrant bisimplicial sets.

**Theorem** ([Rezk](#)).  $(\infty, 1)$ -categories are modeled by [Rezk spaces](#) aka complete Segal spaces.

# Shapes in the theory of the directed interval



Our types may depend on other types and also on **shapes**  $\Phi \subset \mathbb{2}^n$ , polytopes embedded in a directed cube, defined in a language

$$\top, \perp, \wedge, \vee, \equiv \quad \text{and} \quad 0, 1, \leq$$

satisfying **intuitionistic logic** and **strict interval** axioms.

$$\Delta^n := \{(t_1, \dots, t_n) : \mathbb{2}^n \mid t_n \leq \dots \leq t_1\} \quad \text{e.g.} \quad \Delta^1 := \mathbb{2}$$

$$\Delta^2 := \left\{ \begin{array}{ccc} & (t,t) & (1,1) \\ & \diagdown & | \\ (0,0) & & (1,t) \\ & \diagup & \\ & (t,0) & (1,0) \end{array} \right.$$

$$\partial\Delta^2 := \{(t_1, t_2) : \mathbb{2}^2 \mid (t_2 \leq t_1) \wedge ((0 = t_2) \vee (t_2 = t_1) \vee (t_1 = 1))\}$$

$$\Lambda_1^2 := \{(t_1, t_2) : \mathbb{2}^2 \mid (t_2 \leq t_1) \wedge ((0 = t_2) \vee (t_1 = 1))\}$$

Because  $\phi \wedge \psi$  implies  $\phi$ , there are **shape inclusions**  $\Lambda_1^2 \subset \partial\Delta^2 \subset \Delta^2$ .

# Extension types



shape inclusion:  $\Phi := \{t \in \mathcal{P}^n \mid \phi\}$  and  $\Psi = \{t \in \mathcal{P}^n \mid \psi\}$  so that  $\phi$  implies  $\psi$ , i.e., so that  $\Phi \subset \Psi$ .

Formation rule for extension types

$$\frac{\Phi \subset \Psi \text{ shape} \quad A \text{ type} \quad a : \Phi \rightarrow A}{\left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \downarrow & \searrow \text{dashed} & \\ \Psi & & \end{array} \right\rangle \text{ type}}$$

A term  $f : \left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \downarrow & \searrow \text{dashed} & \\ \Psi & & \end{array} \right\rangle$  defines

$$f : \Psi \rightarrow A \text{ so that } f(t) \equiv a(t) \text{ for } t : \Phi.$$

The simplicial type theory allows us to *prove* equivalences between extension types along composites or products of shape inclusions.



3

Segal types and Rezk types

# Hom types

Formation rule for extension types

$$\frac{\Phi \subset \Psi \text{ shape} \quad \Psi \vdash A \text{ type} \quad a : \Phi \rightarrow A}{\left\langle \begin{array}{ccc} \Phi & \xrightarrow{a} & A \\ \Downarrow & \searrow \text{dashed} & \uparrow \\ \Psi & & \end{array} \right\rangle \text{ type}}$$

The **hom type** for  $A$  depends on two terms in  $A$ :

$$x, y : A \vdash \text{hom}_A(x, y)$$

$$\frac{\partial\Delta^1 \subset \Delta^1 \text{ shape} \quad A \text{ type} \quad [x, y] : \partial\Delta^1 \rightarrow A}{\text{hom}_A(x, y) := \left\langle \begin{array}{ccc} \partial\Delta^1 & \xrightarrow{[x, y]} & A \\ \Downarrow & \searrow \text{dashed} & \uparrow \\ \Delta^1 & & \end{array} \right\rangle \text{ type}}$$

A term  $f : \text{hom}_A(x, y)$  defines an **arrow** in  $A$  from  $x$  to  $y$ .

# Segal types have unique binary composites



A type  $A$  is *Segal* iff every composable pair of arrows has a unique composite, i.e., for every  $f : \text{hom}_A(x, y)$  and  $g : \text{hom}_A(y, z)$  the type

$$\left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[f,g]} & A \\ \Downarrow & \dashrightarrow & \\ \Delta^2 & & \end{array} \right\rangle \quad \text{is contractible.}$$

**Prop.** A Reedy fibrant bisimplicial set  $A$  is *Segal* if and only if  $A^{\Delta^2} \rightarrow A^{\Lambda_1^2}$  has contractible fibers.

**Notation.** Let  $\text{comp}_{g,f} : \left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[f,g]} & A \\ \Downarrow & \dashrightarrow & \\ \Delta^2 & & \end{array} \right\rangle$  denote the unique

inhabitant and write  $g \circ f : \text{hom}_A(x, z)$  for its inner face, the composite of  $f$  and  $g$ .

# Identity arrows



For any  $x : A$ , the constant function defines a term

$$\text{id}_x := \lambda t.x : \text{hom}_A(x, x) := \left\langle \begin{array}{ccc} \partial\Delta^1 & \xrightarrow{[x,x]} & A \\ \Downarrow & \nearrow & \\ \Delta^1 & & \end{array} \right\rangle,$$

which we denote by  $\text{id}_x$  and call the **identity arrow**.

For any  $f : \text{hom}_A(x, y)$  in a Segal type  $A$ , the term

$$\lambda(s, t).f(t) : \left\langle \begin{array}{ccc} \Lambda_1^2 & \xrightarrow{[\text{id}_x, f]} & A \\ \Downarrow & \nearrow & \\ \Delta^2 & & \end{array} \right\rangle$$

witnesses the unit axiom  $f = f \circ \text{id}_x$ .

# Associativity of composition




Let  $A$  be a Segal type with arrows

$$f : \text{hom}_A(x, y), \quad g : \text{hom}_A(y, z), \quad h : \text{hom}_A(z, w).$$

Prop.  $h \circ (g \circ f) = (h \circ g) \circ f.$

Proof: Consider the composable arrows in the Segal type  $\Delta^1 \rightarrow A$ :



Composing defines a term in the type  $\Delta^2 \rightarrow (\Delta^1 \rightarrow A)$  which yields a term  $l : \text{hom}_A(x, w)$  so that  $l = h \circ (g \circ f)$  and  $l = (h \circ g) \circ f.$



# Isomorphisms



An arrow  $f: \text{hom}_A(x, y)$  in a Segal type is an **isomorphism** if it has a two-sided inverse  $g: \text{hom}_A(y, x)$ . However, the type

$$\sum_{g: \text{hom}_A(y, x)} (g \circ f = \text{id}_x) \times (f \circ g = \text{id}_y)$$

has higher-dimensional structure and is *not* a **proposition**. Instead define

$$\text{isom}(f) := \left( \sum_{g: \text{hom}_A(y, x)} g \circ f = \text{id}_x \right) \times \left( \sum_{h: \text{hom}_A(y, x)} f \circ h = \text{id}_y \right).$$

For  $x, y : A$ , the **type of isomorphisms** from  $x$  to  $y$  is:

$$x \cong_A y := \sum_{f: \text{hom}_A(x, y)} \text{isom}(f).$$

# Rezk types



By path induction, to define a map

$$\text{id-to-iso} : (x =_A y) \rightarrow (x \cong_A y)$$

for all  $x, y : A$  it suffices to define

$$\text{id-to-iso}(\text{refl}_x) := \text{id}_x.$$

A Segal type  $A$  is **Rezk** if every isomorphism is an identity, i.e., if the map

$$\text{id-to-iso} : (x =_A y) \rightarrow (x \cong_A y)$$

is an equivalence.

# Discrete types



Similarly by path induction define

$$\text{id-to-arr}: \prod_{x,y:A} (x =_A y) \rightarrow \text{hom}_A(x,y) \quad \text{by} \quad \text{id-to-arr}(\text{refl}_x) := \text{id}_x,$$

and call a type  $A$  **discrete** if **id-to-arr** is an equivalence.

**Prop.** A type is discrete if and only if it is Rezk and all of its arrows are isomorphisms. Thus, if the Rezk types are  $(\infty, 1)$ -categories, then the discrete types are  $\infty$ -groupoids.

Proof:

$$\begin{array}{ccc} x =_A y & \xrightarrow{\text{id-to-arr}} & \text{hom}_A(x,y) \\ & \searrow \text{id-to-iso} & \nearrow \\ & x \cong_A y & \end{array}$$



4

The synthetic theory of  
 $(\infty, 1)$ -categories

# Covariant fibrations I



A type family  $x : A \vdash B(x)$  over a Segal type  $A$  is **covariant** if for every  $f : \text{hom}_A(x, y)$  and  $u : B(x)$  there is a unique lift of  $f$  with domain  $u$ , i.e., if

$$\sum_{v:B(y)} \text{hom}_{B(f)}(u, v) \quad \text{is contractible.}$$

Here

$$\text{hom}_{B(f)}(u, v) := \left\langle \begin{array}{ccc} & & B(f) \\ & \nearrow [u,v] & \downarrow \uparrow \\ \partial\Delta^1 & \xrightarrow{\quad} & \Delta^1 \end{array} \right\rangle \quad \text{where} \quad \begin{array}{ccc} B(f) & \longrightarrow & B \\ \downarrow & \lrcorner & \downarrow \\ \Delta^1 & \xrightarrow{f} & A \end{array}$$

is the type of **arrows** in  $B$  from  $u$  to  $v$  **over**  $f$ .

**Notation.** The codomain of the unique lift defines a term  $f_*u : B(y)$ .

**Prop.** For  $u : B(x)$ ,  $f : \text{hom}_A(x, y)$ , and  $g : \text{hom}_A(y, z)$ ,

$$g_*(f_*u) = (g \circ f)_*u \quad \text{and} \quad (\text{id}_x)_*u = u.$$



A type family  $x : A \vdash B(x)$  over a Segal type  $A$  is **covariant** if for every  $f : \text{hom}_A(x, y)$  and  $u : B(x)$  there is a unique lift of  $f$  with domain  $u$ .

**Prop.** If  $x : A \vdash B(x)$  is covariant then for each  $x : A$  the fiber  $B(x)$  is discrete. Thus covariant type families are fibered in  $\infty$ -groupoids.

**Prop.** Fix  $a : A$ . The type family  $x : A \vdash \text{hom}_A(a, x)$  is covariant.

For  $u : \text{hom}_A(a, x)$  and  $f : \text{hom}_A(x, y)$ , the transport  $f_*u$  equals the composite  $f \circ u$  as terms in  $\text{hom}_A(a, y)$ , i.e.,  $f_*(u) = f \circ u$ .

# The Yoneda lemma



Let  $x : A \vdash B(x)$  be a covariant family over a Segal type and fix  $a : A$ .

Yoneda lemma. The maps

$$\text{ev-id} := \lambda\phi.\phi(a, \text{id}_a) : \left( \prod_{x:A} \text{hom}_A(a, x) \rightarrow B(x) \right) \rightarrow B(a)$$

and

$$\text{yon} := \lambda u.\lambda x.\lambda f.f_*u : B(a) \rightarrow \left( \prod_{x:A} \text{hom}_A(a, x) \rightarrow B(x) \right)$$

are inverse equivalences.

**Proof:** The transport operation for covariant families is functorial in  $A$  and fiberwise maps between covariant families are automatically natural.

**Note.** A representable isomorphism  $\phi : \prod_{x:A} \text{hom}_A(a, x) \cong \text{hom}_A(b, x)$  induces an identity  $\text{ev-id}(\phi) : b =_A a$  if the Segal type  $A$  is Rezk.

# The dependent Yoneda lemma



*From a type-theoretic perspective, the Yoneda lemma is a “directed” version of the “transport” operation for identity types. This suggests a “dependently typed” generalization of the Yoneda lemma, analogous to the full induction principle for identity types.*

**Dependent Yoneda lemma.** If  $A$  is a Segal type and  $B(x, y, f)$  is a covariant family dependent on  $x, y : A$  and  $f : \text{hom}_A(x, y)$ , then evaluation at  $(x, x, \text{id}_x)$  defines an equivalence

$$\text{ev-id} : \left( \prod_{x, y : A} \prod_{f : \text{hom}_A(x, y)} B(x, y, f) \right) \rightarrow \prod_{x : A} B(x, x, \text{id}_x)$$

This is useful for proving equivalences between various types of coherent or incoherent adjunction data.



# Dependent Yoneda is directed path induction



Takeaway: the dependent Yoneda lemma is directed path induction.

**Path induction:** If  $B(x, y, p)$  is a type family dependent on  $x, y : A$  and  $p : x =_A y$ , then there is a function

$$\text{path-ind} : \left( \prod_{x:A} B(x, x, \text{refl}_x) \right) \rightarrow \left( \prod_{x,y:A} \prod_{p:x=_A y} B(x, y, p) \right).$$

Thus, to prove  $B(x, y, p)$  it suffices to assume  $y$  is  $x$  and  $p$  is  $\text{refl}_x$ .

**Dependent Yoneda Lemma:** If  $B(x, y, f)$  is a covariant family dependent on  $x, y : A$  and  $f : \text{hom}_A(x, y)$  and  $A$  is Segal, then there is a function

$$\text{id-ind} : \left( \prod_{x:A} B(x, x, \text{id}_x) \right) \rightarrow \left( \prod_{x,y:A} \prod_{f:\text{hom}_A(x,y)} B(x, y, f) \right).$$

Thus, to prove  $B(x, y, f)$  it suffices to assume  $y$  is  $x$  and  $f$  is  $\text{id}_x$ .

# References



For considerably more, see:

Emily Riehl and Michael Shulman, [A type theory for synthetic  \$\infty\$ -categories](#), [arXiv:1705.07442](#)

To explore homotopy type theory:

[Homotopy Type Theory: Univalent Foundations of Mathematics](#),  
<https://homotopytypetheory.org/book/>

Michael Shulman, [Homotopy type theory: the logic of space](#),  
[arXiv:1703.03007](#)

Thank you!